

Polytopes for Crystallized Demazure Modules and Extremal Vectors

Toshiki NAKASHIMA

Department of Mathematics,
Sophia University, Tokyo 102-8554, JAPAN
e-mail: toshiki@mm.sophia.ac.jp

1 Introduction

Demazure's character formula for arbitrary Kac-Moody Lie algebra was given by S.Kumar and O.Mathieu independently ([6],[8]) by using geometric methods. In 1995, P.Littelmann gave some conjecture (partially solved by himself) about the relation between Demazure's character formula and crystal bases [7], which was solved affirmatively by M.Kashiwara [3]. Then it gave purely algebraic proof for Demazure's character formula for symmetrizable Kac-Moody Lie algebras. Here let us see those formulations. Let $\mathfrak{g}$ be a symmetrizable Kac-Moody Lie algebra (in the context of "crystal base", we need "symmetrizable"), and $\mathfrak{n}^+$ be the nilpotent subalgebra of $\mathfrak{g}$. Furthermore, let $\mathbf{Z}[P]$ be the group algebra of the weight lattice P and W be the Weyl group associated with $\mathfrak{g}$. Then Demazure operator $D_w : \mathbf{Z}[P] \rightarrow \mathbf{Z}[P]$ ($w \in W$) is given as follows: for $i \in I$ (index set) we set $D_i(e^\lambda) := e^\lambda(1 - e^{-(1+\langle h_i, \lambda \rangle)\alpha_i})/(1 - e^{-\alpha_i})$ and for $w = s_{i_1} \cdots s_{i_l}$ set $D_w := D_{i_1} \cdots D_{i_l}$, which is well-defined. Let $V(\lambda)$ be the irreducible highest weight module with the highest weight λ and $u_{w\lambda}$ be the extremal vector with the weight $w\lambda$ ($w \in W$). Then, Demazure's character formula is described as follows:

$$ch(U(\mathfrak{n}^+)u_{w\lambda}) = D_w(e^\lambda). \quad (1.1)$$

In [7], Littelmann gave the following conjecture: Let $V(\lambda)$ be the irreducible $U_q(\mathfrak{g})$ -highest weight module with the highest weight λ and $(L(\lambda), B(\lambda))$ be its crystal base. Then there exists a subset $B_w(\lambda) \subset B(\lambda)$ such that

$$U_q^+(\mathfrak{g})u_{w\lambda} \cap L(\lambda)/U_q^+(\mathfrak{g})u_{w\lambda} \cap qL(\lambda) = \bigoplus_{b \in B_w(\lambda)} \mathbb{Q}b, \quad (1.2)$$

$$\sum_{b \in B_w(\lambda)} b = \mathfrak{D}_{i_1} \cdots \mathfrak{D}_{i_l} u_\lambda, \quad (1.3)$$

where u_λ is the highest weight vector with the weight λ and $\mathfrak{D}_i$ is the additive operator on $\mathbf{Z}^{\oplus B(\lambda)}$ given by:

$$\mathfrak{D}_i b := \begin{cases} \sum_{0 \leq k \leq \langle h_i, wt(b) \rangle} \tilde{f}_i^k b & \text{if } \langle h_i, wt(b) \rangle \geq 0, \\ -\sum_{1 \leq k < -\langle h_i, wt(b) \rangle} \tilde{e}_i^k b & \text{if } \langle h_i, wt(b) \rangle < 0. \end{cases}$$

We call the left-hand side of (1.2) *crystallized Demazure module* of $V(\lambda)$ associated with $w \in W$. Here we know that Littelmann's conjecture implies Demazure's character formula by the following way: Define the operator $ewt :$

$\mathbf{Z}^{\oplus B(\lambda)} \longrightarrow \mathbf{Z}[P]$ by $ewt(b) := e^{wt(b)}$ for $b \in B(\lambda)$ and $ewt(b_1 + b_2) = ewt(b_1) + ewt(b_2)$. Now, we have $ewt(\mathfrak{D}_i b) = D_i(ewt(b))$. Thus, by (1.2) and (1.3) we have

$$\begin{aligned} ch(U_q^+(\mathfrak{g})u_{w\lambda}) &= ewt(\sum_{b \in B_w(\lambda)} b) = ewt(\mathfrak{D}_{i_l} \cdots \mathfrak{D}_{i_1} u_\lambda) \\ &= D_{i_l} \cdots D_{i_1} ewt(u_\lambda) = D_{i_l} \cdots D_{i_1}(e^\lambda) = D_w(e^\lambda). \end{aligned}$$

In [3], Kashiwara shown the existence of $B_w(\lambda)$ for arbitrary symmetrizable Kac-Moody cases and characterized it as follows:

Theorem 1.1 ([3]) (i) $\tilde{e}_i B_w(\lambda) \subset B_w(\lambda) \sqcup \{0\}$.

(ii) If $s_i w < w$ (Bruhat order), then $B_w(\lambda) = \{\tilde{f}_i^k b; k \geq 0, b \in B_{s_i w}(\lambda), \tilde{e}_i b = 0\} \setminus \{0\}$.

(iii) For any i -string S , $S \cap B_w(\lambda)$ is either empty or S or $\{the\ highest\ weight\ vector\ of\ S\}$.

In [9],[10], we developed the polyhedral realization of crystal bases. We shall explain the relations between crystal bases of Demazure modules and the polyhedral realizations briefly. Let $\iota = \cdots i_k, \cdots, i_2, i_1$ be an infinite sequence from the index set I satisfying some condition and λ be a dominant integral weight. Then there exists the embedding $\Psi_t^{(\lambda)} : B(\lambda) \hookrightarrow \mathbf{Z}_t^\infty[\lambda] (\cong \mathbf{Z}^\infty)$. The exact image of $\Psi_t^{(\lambda)}$ is described (under some assumption) as a subset in $\mathbf{Z}^\infty$ given by some system of linear inequalities, which is called *polyhedral realization*. Let $w = s_{i_l} \cdots s_{i_1}$ (reduced expression) be an element in W and take a sequence $\iota = (j_k)_{k \geq 1}$ which satisfies $i_k = j_k$ ($1 \leq k \leq l$). Then in this paper, the subset $\Psi_t^{(\lambda)}(B_w(\lambda))$ is given as a set of lattice points of some convex polytope in $\mathbf{Z}^\infty$, where “polytope” means a bounded polyhedron. Furthermore, we succeed in giving explicit form of extremal vector $\Psi_t^{(\lambda)}(u_{w\lambda})$ which is contained in $\Psi_t^{(\lambda)}(B_w(\lambda))$ as the unique solution of some system of linear equations.

The organization of this paper is as follows: In Sect.2 we review the polyhedral realizations of crystals. We shall describe the polytopes for $B_w(\lambda)$ in Sect. 3 and the extremal vectors in Sect.4.

2 Polyhedral realizations of crystals

2.1 Notations

We list the notations used in this paper. Most of them are same as those in [10].

Let $\mathfrak{g}$ be a symmetrizable Kac-Moody algebra over $\mathbf{Q}$ with a Cartan subalgebra $\mathfrak{t}$, a weight lattice $P \subset \mathfrak{t}^*$, the set of simple roots $\{\alpha_i : i \in I\} \subset \mathfrak{t}^*$, and the set of coroots $\{h_i : i \in I\} \subset \mathfrak{t}$, where I is a finite index set. Let $\langle h, \lambda \rangle$ be the pairing between $\mathfrak{t}$ and $\mathfrak{t}^*$, and (α, β) be an inner product on $\mathfrak{t}^*$ such that $(\alpha_i, \alpha_i) \in 2\mathbf{Z}_{\geq 0}$ and $\langle h_i, \lambda \rangle = \frac{2(\alpha_i, \lambda)}{(\alpha_i, \alpha_i)}$ for $\lambda \in \mathfrak{t}^*$. Let $P^* = \{h \in \mathfrak{t} : \langle h, P \rangle \subset \mathbf{Z}\}$ and $P_+ := \{\lambda \in P : \langle h_i, \lambda \rangle \in \mathbf{Z}_{\geq 0}\}$. We call an element in P_+ a *dominant integral weight*. Here we define a partial order on P by: For $\lambda, \mu \in P$, $\lambda \succ \mu \Leftrightarrow \lambda - \mu \in \oplus_{i \in I} \mathbf{Q}_{\geq 0} \alpha_i$. The quantum algebra $U_q(\mathfrak{g})$ is an associative $\mathbf{Q}(q)$ -algebra generated by the e_i, f_i ($i \in I$), and q^h ($h \in P^*$) satisfying the usual relations. The algebra $U_q^-(\mathfrak{g})$ is the subalgebra of $U_q(\mathfrak{g})$ generated by the f_i ($i \in I$).

For the irreducible highest weight module of $U_q(\mathfrak{g})$ with the highest weight $\lambda \in P_+$, we denote $V(\lambda)$ and its *crystal base* we denote $(L(\lambda), B(\lambda))$. Similarly, for the crystal base of the algebra $U_q^-(\mathfrak{g})$ we denote $(L(\infty), B(\infty))$ (see [1],[2],[4]). Let $\pi_\lambda : U_q^-(\mathfrak{g}) \longrightarrow V(\lambda) \cong U_q^-(\mathfrak{g})/\sum_i U_q^-(\mathfrak{g})\tilde{f}_i^{1+\langle h_i, \lambda \rangle}$ be the canonical projection and $\hat{\pi}_\lambda : L(\infty)/qL(\infty) \longrightarrow L(\lambda)/qL(\lambda)$ be the induced map from π_λ . Here note that $\hat{\pi}_\lambda(B(\infty)) = B(\lambda) \sqcup \{0\}$.

By the terminology *crystal* we mean some combinatorial object obtained by abstracting the properties of crystal bases. Indeed, crystal constitutes a set B and the maps $wt : B \longrightarrow P$, $\varepsilon_i, \varphi_i : B \longrightarrow \mathbf{Z} \sqcup \{-\infty\}$ and $\tilde{e}_i, \tilde{f}_i : B \sqcup \{0\} \longrightarrow B \sqcup \{0\}$ ($i \in I$) with several axioms (see [3],[9],[10]). In fact, $B(\infty)$ and $B(\lambda)$ are the typical examples of crystals.

It is well-known that $U_q(\mathfrak{g})$ has a Hopf algebra structure. Then the tensor product of $U_q(\mathfrak{g})$ -modules has a $U_q(\mathfrak{g})$ -module structure. The crystal bases have very nice properties for tensor operations. Indeed, if (L_i, B_i) is a crystal base of $U_q(\mathfrak{g})$ -module M_i ($i = 1, 2$), $(L_1 \otimes_A L_2, B_1 \otimes B_2)$ is a crystal base of $M_1 \otimes_{\mathbf{Q}(q)} M_2$ ([2]). Consequently, we can consider the tensor product of crystals and then they constitute a tensor category.

2.2 Polyhedral Realization of $B(\infty)$

In this subsection, we recall the results in [9].

Consider the additive group

$$\mathbf{Z}^\infty := \{(\cdots, x_k, \cdots, x_2, x_1) : x_k \in \mathbf{Z} \text{ and } x_k = 0 \text{ for } k \gg 0\}; \quad (2.1)$$

we will denote by $\mathbf{Z}_{\geq 0}^\infty \subset \mathbf{Z}^\infty$ the subsemigroup of nonnegative sequences. To the rest of this section, we fix an infinite sequence of indices $\iota = \cdots, i_k, \cdots, i_2, i_1$ from I such that

$$i_k \neq i_{k+1} \text{ and } \#\{k : i_k = i\} = \infty \text{ for any } i \in I. \quad (2.2)$$

We can associate to ι a crystal structure on $\mathbf{Z}^\infty$ and denote it by $\mathbf{Z}_\iota^\infty$ ([9, 2.4]).

Proposition 2.1 ([3], See also [9]) *There is a unique embedding of crystals (called Kashiwara embedding)*

$$\Psi_\iota : B(\infty) \hookrightarrow \mathbf{Z}_{\geq 0}^\infty \subset \mathbf{Z}_\iota^\infty, \quad (2.3)$$

such that $\Psi_\iota(u_\infty) = (\cdots, 0, \cdots, 0, 0)$.

Consider the infinite dimensional vector space

$$\mathbf{Q}^\infty := \{x = (\cdots, x_k, \cdots, x_2, x_1) : x_k \in \mathbf{Q} \text{ and } x_k = 0 \text{ for } k \gg 0\},$$

and its dual space $(\mathbf{Q}^\infty)^* := \text{Hom}(\mathbf{Q}^\infty, \mathbf{Q})$. We will write a linear form $\varphi \in (\mathbf{Q}^\infty)^*$ as $\varphi(x) = \sum_{k \geq 1} \varphi_k x_k$ ($\varphi_j \in \mathbf{Q}$).

For the fixed infinite sequence $\iota = (i_k)$ we set $k^{(+)} := \min\{l : l > k \text{ and } i_k = i_l\}$ and $k^{(-)} := \max\{l : l < k \text{ and } i_k = i_l\}$ if it exists, or $k^{(-)} = 0$ otherwise. We set for $x \in \mathbf{Q}^\infty$, $\beta_0(x) = 0$ and

$$\beta_k(x) := x_k + \sum_{k < j < k^{(+)}} \langle h_{i_k}, \alpha_{i_j} \rangle x_j + x_{k^{(+)}} \quad (k \geq 1). \quad (2.4)$$

We define a piecewise-linear operator $S_k = S_{k,\iota}$ on $(\mathbf{Q}^\infty)^*$ by

$$S_k(\varphi) := \begin{cases} \varphi - \varphi_k \beta_k & \text{if } \varphi_k > 0, \\ \varphi - \varphi_k \beta_{k^{(-)}} & \text{if } \varphi_k \leq 0. \end{cases}$$

Here we set

$$\Xi_\iota := \{S_{j_l} \cdots S_{j_2} S_{j_1} x_{j_0} \mid l \geq 0, j_0, j_1, \dots, j_l \geq 1\}, \quad (2.5)$$

$$\Sigma_\iota := \{x \in \mathbf{Z}^\infty \subset \mathbf{Q}^\infty \mid \varphi(x) \geq 0 \text{ for any } \varphi \in \Xi_\iota\}. \quad (2.6)$$

We impose on ι the following positivity assumption:

$$\text{if } k^{(-)} = 0 \text{ then } \varphi_k \geq 0 \text{ for any } \varphi(x) = \sum_k \varphi_k x_k \in \Xi_\iota. \quad (2.7)$$

Theorem 2.1 ([9]) *Let ι be a sequence of indices satisfying (2.2) and (2.7). Then we have $\text{Im}(\Psi_\iota)(\cong B(\infty)) = \Sigma_\iota$.*

2.3 Polyhedral Realization of $B(\lambda)$

In this subsection, we review the result in [10]. In the rest of this section, λ is supposed to be a dominant integral weight. Let $R_\lambda := \{r_\lambda\}$ be the crystal defined in [10]. Consider the crystal $B(\infty) \otimes R_\lambda$ and define the map

$$\Phi_\lambda : (B(\infty) \otimes R_\lambda) \sqcup \{0\} \longrightarrow B(\lambda) \sqcup \{0\}, \quad (2.8)$$

by $\Phi_\lambda(0) = 0$ and $\Phi_\lambda(b \otimes r_\lambda) = \hat{\pi}_\lambda(b)$ for $b \in B(\infty)$. We set

$$\tilde{B}(\lambda) := \{b \otimes r_\lambda \in B(\infty) \otimes R_\lambda \mid \Phi_\lambda(b \otimes r_\lambda) \neq 0\}.$$

Theorem 2.2 ([10]) (i) *The map Φ_λ becomes a surjective strict morphism of crystals $B(\infty) \otimes R_\lambda \longrightarrow B(\lambda)$.*

(ii) *$\tilde{B}(\lambda)$ is a subcrystal of $B(\infty) \otimes R_\lambda$, and Φ_λ induces the isomorphism of crystals $\tilde{B}(\lambda) \xrightarrow{\sim} B(\lambda)$.*

Let us denote $\mathbf{Z}_\iota^\infty \otimes R_\lambda$ by $\mathbf{Z}_\iota^\infty[\lambda]$. Here note that since the crystal R_λ has only one element, as a set we can identify $\mathbf{Z}_\iota^\infty[\lambda]$ with $\mathbf{Z}_\iota^\infty$ but their crystal structures are different. As for the explicit crystal structure of $\mathbf{Z}_\iota^\infty[\lambda]$, see 3.1 below. By Theorem 2.2, we have the strict embedding of crystals $\Omega_\lambda : B(\lambda)(\cong \tilde{B}(\lambda)) \hookrightarrow B(\infty) \otimes R_\lambda$. Combining Ω_λ and the Kashiwara embedding Ψ_ι , we obtain the following:

Theorem 2.3 ([10]) *There exists the unique strict embedding of crystals*

$$\Psi_\iota^{(\lambda)} : B(\lambda) \xrightarrow{\Omega_\lambda} B(\infty) \otimes R_\lambda \xrightarrow{\Psi_\iota \otimes \text{id}} \mathbf{Z}_\iota^\infty \otimes R_\lambda =: \mathbf{Z}_\iota^\infty[\lambda], \quad (2.9)$$

such that $\Psi_\iota^{(\lambda)}(u_\lambda) = (\cdots, 0, 0, 0) \otimes r_\lambda$.

We fix a sequence of indices ι satisfying (2.2) and take a dominant integral weight $\lambda \in P_+$. For $k \geq 1$ let $k^{(\pm)}$ be the ones in 2.2. Let $\beta_k^{(\pm)}(x)$ be linear functions given by

$$\beta_k^{(+)}(x) = \sigma_k(x) - \sigma_{k^{(+)}}(x) = x_k + \sum_{k < j < k^{(+)}} \langle h_{i_k}, \alpha_{i_j} \rangle x_j + x_{k^{(+)}}, \quad (2.10)$$

$$\begin{aligned} & \beta_k^{(-)}(x) \\ &= \begin{cases} \sigma_{k^{(-)}}(x) - \sigma_k(x) = x_{k^{(-)}} + \sum_{k^{(-)} < j < k} \langle h_{i_k}, \alpha_{i_j} \rangle x_j + x_k & \text{if } k^{(-)} > 0, \\ \sigma_0^{(i_k)}(x) - \sigma_k(x) = -\langle h_{i_k}, \lambda \rangle + \sum_{1 \leq j < k} \langle h_{i_k}, \alpha_{i_j} \rangle x_j + x_k & \text{if } k^{(-)} = 0, \end{cases} \end{aligned} \quad (2.11)$$

(As for the functions σ_k and $\sigma_0^{(i)}$ see (3.1) and (3.2) below.). Here note that $\beta_k^{(+)} = \beta_k$ and $\beta_k^{(-)} = \beta_{k^{(-)}}$ if $k^{(-)} > 0$.

Using this notation, for every $k \geq 1$, we define an operator $\widehat{S}_k = \widehat{S}_{k,\iota}$ for a linear function $\varphi(x) = c + \sum_{k \geq 1} \varphi_k x_k$ ($c, \varphi_k \in \mathbf{Q}$) on $\mathbf{Q}^\infty$ by:

$$\widehat{S}_k(\varphi) := \begin{cases} \varphi - \varphi_k \beta_k^{(+)} & \text{if } \varphi_k > 0, \\ \varphi - \varphi_k \beta_k^{(-)} & \text{if } \varphi_k \leq 0. \end{cases}$$

For the fixed sequence $\iota = (i_k)$, in case $k^{(-)} = 0$ for $k \geq 1$, there exists unique $i \in I$ such that $i_k = i$. We denote such k by $\iota^{(i)}$, namely, $\iota^{(i)}$ is the first number k such that $i_k = i$. Here for $\lambda \in P_+$ and $i \in I$ we set

$$\lambda^{(i)}(x) := -\beta_{\iota^{(i)}}^{(-)}(x) = \langle h_i, \lambda \rangle - \sum_{1 \leq j < \iota^{(i)}} \langle h_i, \alpha_{i_j} \rangle x_j - x_{\iota^{(i)}}. \quad (2.12)$$

For ι and a dominant integral weight λ , let $\Xi_\iota[\lambda]$ be the set of all linear functions generated by $\widehat{S}_k = \widehat{S}_{k,\iota}$ from the functions x_j ($j \geq 1$) and $\lambda^{(i)}$ ($i \in I$), namely,

$$\begin{aligned} \Xi_\iota[\lambda] &:= \{ \widehat{S}_{j_l} \cdots \widehat{S}_{j_1} x_{j_0} : l \geq 0, j_0, \dots, j_l \geq 1 \} \\ &\cup \{ \widehat{S}_{j_k} \cdots \widehat{S}_{j_1} \lambda^{(i)}(x) : k \geq 0, i \in I, j_1, \dots, j_k \geq 1 \}. \end{aligned} \quad (2.13)$$

Now we set

$$\Sigma_\iota[\lambda] := \{x \in \mathbf{Z}_\iota^\infty[\lambda] \subset \mathbf{Q}^\infty : \varphi(x) \geq 0 \text{ for any } \varphi \in \Xi_\iota[\lambda]\}. \quad (2.14)$$

For a sequence ι and a dominant integral weight λ , a pair (ι, λ) is called *ample* if $\Sigma_\iota[\lambda] \ni \vec{0} = (\dots, 0, 0)$.

Theorem 2.4 ([10]) *Suppose that (ι, λ) is ample. Then we have $\text{Im}(\Psi_\iota^{(\lambda)}) (\cong B(\lambda)) = \Sigma_\iota[\lambda]$.*

3 Crystallized Demazure modules

3.1 Structure of $\mathbf{Z}_\iota^\infty[\lambda]$

We shall review an explicit crystal structure of $\mathbf{Z}_\iota^\infty[\lambda]$ in [10]. Fix a sequence of indices $\iota := (i_k)_{k \geq 1}$ satisfying the condition (2.2) and a weight $\lambda \in P$. (Here

we do not necessarily assume that λ is dominant.) As we stated before, we can identify $\mathbf{Z}^\infty$ with $\mathbf{Z}^\infty[\lambda]$ as a set. Thus $\mathbf{Z}^\infty[\lambda]$ can be regarded as a subset of $\mathbf{Q}^\infty$, and then we denote an element in $\mathbf{Z}^\infty[\lambda]$ by $x = (\cdots, x_k, \cdots, x_2, x_1)$. For $x = (\cdots, x_k, \cdots, x_2, x_1) \in \mathbf{Q}^\infty$ we define the linear functions

$$\sigma_k(x) := x_k + \sum_{j>k} \langle h_{i_k}, \alpha_{i_j} \rangle x_j, \quad (k \geq 1) \quad (3.1)$$

$$\sigma_0^{(i)}(x) := -\langle h_i, \lambda \rangle + \sum_{j \geq 1} \langle h_i, \alpha_{i_j} \rangle x_j, \quad (i \in I) \quad (3.2)$$

Here note that since $x_j = 0$ for $j \gg 0$ on $\mathbf{Q}^\infty$, the functions σ_k and $\sigma_0^{(i)}$ are well-defined. Let $\sigma^{(i)}(x) := \max_{k: i_k = i} \sigma_k(x)$, and $M^{(i)} := \{k : i_k = i, \sigma_k(x) = \sigma^{(i)}(x)\}$. Note that $\sigma^{(i)}(x) \geq 0$, and that $M^{(i)} = M^{(i)}(x)$ is a finite set if and only if $\sigma^{(i)}(x) > 0$. Now we define the maps $\tilde{e}_i : \mathbf{Z}^\infty[\lambda] \sqcup \{0\} \longrightarrow \mathbf{Z}^\infty[\lambda] \sqcup \{0\}$ and $\tilde{f}_i : \mathbf{Z}^\infty[\lambda] \sqcup \{0\} \longrightarrow \mathbf{Z}^\infty[\lambda] \sqcup \{0\}$ by setting $\tilde{e}_i(0) = \tilde{f}_i(0) = 0$, and

$$(\tilde{f}_i(x))_k = x_k + \delta_{k, \min M^{(i)}} \text{ if } \sigma^{(i)}(x) > \sigma_0^{(i)}(x); \text{ otherwise } \tilde{f}_i(x) = 0, \quad (3.3)$$

$$(\tilde{e}_i(x))_k = x_k - \delta_{k, \max M^{(i)}} \text{ if } \sigma^{(i)}(x) > 0 \text{ and } \sigma^{(i)}(x) \geq \sigma_0^{(i)}(x); \text{ otherwise } \tilde{e}_i(x) = 0, \quad (3.4)$$

where $\delta_{i,j}$ is the Kronecker's delta. We also define the functions wt , ε_i and φ_i on $\mathbf{Z}^\infty[\lambda]$ by

$$wt(x) := \lambda - \sum_{j=1}^{\infty} x_j \alpha_{i_j}, \quad (3.5)$$

$$\varepsilon_i(x) := \max(\sigma^{(i)}(x), \sigma_0^{(i)}(x)) \quad (3.6)$$

$$\varphi_i(x) := \langle h_i, wt(x) \rangle + \varepsilon_i(x). \quad (3.7)$$

Note that by (3.5) we have

$$\langle h_i, wt(x) \rangle = -\sigma_0^{(i)}(x). \quad (3.8)$$

3.2 Polytopes for $B_w(\lambda)$

In this section, we describe the explicit form of the polytopes corresponding to the crystals of Demazure module $B_w(\lambda)$ ($\lambda \in P_+$) as in the introduction.

By the characterization of $B_w(\lambda)$ given in Theorem 1.1 (ii), we can construct it inductively according to some reduced expression of w . Indeed, we have $B_1(\lambda) = \{u_\lambda\}$ (where 1 is the identity of W) and then we obtain $B_{s_i}(\lambda) = \{\tilde{f}_i^k u_\lambda; k \geq 0\} \setminus \{0\}$. Since $\vec{0} := (\cdots, 0, 0)$ corresponds to the highest weight vector, by (3.3) the image by $\Psi_\iota^{(\lambda)}$ is given by;

$$\Sigma_{s_i}[\lambda] := \{(\cdots, 0, 0, k); 0 \leq k \leq \langle h_i, \lambda \rangle\},$$

where $i_1 = i$ for $\iota = (i_k)_{k \geq 1}$.

For $w \in W$, let us fix one reduced expression $w = s_{i_L} s_{i_{L-1}} \cdots s_{i_2} s_{i_1}$ and let $\iota := (j_k)_{k \geq 1}$ be the infinite sequence of indices such that $i_k = j_k$ for $1 \leq k \leq L$. Here we do not necessarily assume that (ι, λ) is ample. In this setting, we have

Proposition 3.1 *Set*

$$\Sigma_w[\lambda] := \{(\cdots, x_k, x_{k-1}, \cdots, x_2, x_1) \in \text{Im}(\Psi_t^{(\lambda)}) \mid x_k = 0 \text{ for } k > L\}. \quad (3.9)$$

Then we have $\Psi_t^{(\lambda)}(B_w(\lambda)) = \Sigma_w[\lambda]$.

Proof. We shall show by induction on the length of w . If the length of w is equal to 0, then $w = 1$. So we have $B_1(\lambda) = \{u_\lambda\}$ and then $\Psi_t(B_1(\lambda)) = \{(\cdots, 0, 0)\}$. If the length of w is equal to 1, then we can set $w = s_{i_1}$. As we have mentioned above, the image of $B_{s_{i_1}}(\lambda)$ by $\Psi_t^{(\lambda)}$ is

$$\{(\cdots, x_2, x_1) \in \text{Im}(\Psi_t^{(\lambda)}) \mid x_2 = x_3 = \cdots = 0\} = \Sigma_{s_{i_1}}[\lambda].$$

Fix $w = s_{i_L} s_{i_{L-1}} \cdots s_{i_2} s_{i_1}$ (reduced expression), and set $w' := s_{i_{L-1}} \cdots s_{i_2} s_{i_1}$. By the hypothesis of the induction, we have

$$\Psi_t^{(\lambda)}(B_{w'}(\lambda)) = \Sigma_{w'}[\lambda] := \{(\cdots, x_k, \cdots, x_2, x_1) \in \text{Im}(\Psi_t^{(\lambda)}) \mid x_k = 0 \text{ for } k > L-1\}. \quad (3.10)$$

Here we show

$$\tilde{f}_{i_L}^l x \in \Sigma_w[\lambda] \cup \{0\} \quad (\text{for any } x \in \Sigma_{w'}[\lambda] \text{ and any } l \in \mathbf{Z}_{\geq 0}), \quad (3.11)$$

by the induction on l . For $x \in \Sigma_{w'}[\lambda]$ and $k > L$ such that $i_k = i_L$, we have $\sigma_k(x) = \sigma_L(x) = 0$ (as for σ_k see (3.1)). It follows from (3.3), that if $\tilde{f}_{i_L} x \neq 0$, then its k -th entry is equal to 0. Thus, we have

$$\tilde{f}_{i_L} x \in \Sigma_w[\lambda] \cup \{0\}.$$

Suppose that

$$\tilde{f}_{i_L}^l x \in \Sigma_w[\lambda] \quad (3.12)$$

and set its L -th entry $x'_L (\geq 0)$. By (3.12), we have $\sigma_k(\tilde{f}_{i_L}^l x) = 0$ ($k > L$ and $i_k = i_L$) and also we have $\sigma_L(\tilde{f}_{i_L}^l x) = x'_L \geq 0$. This implies

$$\sigma_k(\tilde{f}_{i_L}^l x) \leq \sigma_L(\tilde{f}_{i_L}^l x) \leq \sigma^{(i_L)}(\tilde{f}_{i_L}^l x). \quad (3.13)$$

It follows from (3.3) again that we have $\tilde{f}_{i_L}^{l+1} x \in \Sigma_w[\lambda] \cup \{0\}$ and then $\Psi_t^{(\lambda)}(B_w(\lambda)) \subset \Sigma_w[\lambda]$.

Next, we are going to show the opposite inclusion. For any $x = (\cdots, x_k, \cdots, x_2, x_1) \in \Sigma_w[\lambda]$, by (3.6) we have

$$\varepsilon_{i_L}(x) = \max_{k; i_k = i_L} \{\sigma_k(x), \sigma_0^{(i_L)}(x)\} \geq \sigma_L(x) = x_L. \quad (3.14)$$

Since the action of $\tilde{e}_i$ only reduces some entry in x , we have $\tilde{e}_{i_L}^{\varepsilon_{i_L}(x)}(x) \in \Sigma_w[\lambda]$, (note that $\tilde{e}_{i_L}^{\varepsilon_{i_L}(x)}(x)$ is never 0) and

$$\varepsilon_{i_L}(\tilde{e}_{i_L}^{\varepsilon_{i_L}(x)}(x)) = 0. \quad (3.15)$$

By (3.14) and (3.15), we have

$$(\tilde{e}_{i_L}^{\varepsilon_{i_L}(x)}(x))_L (= L\text{-th entry of } \tilde{e}_{i_L}^{\varepsilon_{i_L}(x)}(x)) = 0. \quad (3.16)$$

Thus, we have $\tilde{e}_{i_L}^{\varepsilon_{i_L}(x)}(x) \in \Sigma_{w'}[\lambda]$. Therefore, by Theorem 1.1(ii), we get

$$x \in \tilde{f}_{i_L}^{\varepsilon_{i_L}(x)} \Sigma_{w'}[\lambda] = \tilde{f}_{i_L}^{\varepsilon_{i_L}} \Psi_\iota^{(\lambda)}(B_{w'}(\lambda)) \subset \Psi_\iota^{(\lambda)}(B_w(\lambda)). \quad (3.17)$$

Now we obtain the opposite inclusion $\Sigma_w[\lambda] \subset \Psi_\iota^{(\lambda)}(B_w(\lambda))$ and then completed the proof. $\square$

Practically, we need the assumption “ample”. If (λ, ι) is ample, we can write Proposition 3.1 in the following form:

Proposition 3.2 *If (λ, ι) is ample, we have*

$$\Psi_\iota^{(\lambda)}(B_w(\lambda)) (= \Sigma_w[\lambda]) = \{x = (x_k) \in \mathbf{Z}_\iota^\infty[\lambda] \mid \begin{array}{l} \varphi(x) \geq 0 \text{ for any } \varphi \in \Xi_\iota[\lambda], \\ x_k = 0 \text{ for } k > L. \end{array} \}, \quad (3.18)$$

where $\Xi_\iota[\lambda]$ is given in (2.13).

Now, we obtain the convex “polytope” for $B_w(\lambda)$.

In [3], Kashiwara also introduced the crystal $B_w(\infty) \subset B(\infty)$. This possesses the following remarkable property:

If $b \in B(\infty)$ and $w \in W$ satisfy $\tilde{f}_i b \in B_w(\infty)$, then $\tilde{f}_i^k b \in B_w(\infty)$ for any $k \geq 0$.

This is used for proving Theorem 1.1 (iii).

It is characterized by the following;

- (i) $B_w(\infty) = \{u_\infty\}$ if $w = 1$.
- (ii) if $s_i w < w$, then $B_w(\infty) = \bigcup_{k \geq 0} \tilde{f}_i^k B_{s_i w}(\infty)$.

This implies that $B_w(\infty)$ has also the similar description to $B_w(\lambda)$.

Proposition 3.3 (i) *We have*

$$\Psi_\iota(B_w(\infty)) = \{(\cdots, x_k, \cdots, x_2, x_1) \in \text{Im}(\Psi_\iota) \mid x_k = 0 \text{ for } k > L\}$$

(ii) *If ι satisfies the condition (2.7), we have*

$$\Psi_\iota(B_w(\infty)) = \left\{ x = (\cdots, x_k, \cdots, x_2, x_1) \in \mathbf{Z}_\iota^\infty \mid \begin{array}{l} \varphi(x) \geq 0 \text{ for any } \varphi \in \Xi_\iota \\ x_k = 0 \text{ for } k > L \end{array} \right\},$$

where Ξ_ι is given in (2.5).

3.3 Semi-simple cases

In this subsection, let $\mathfrak{g}$ be a semi-simple Lie algebra, W be the corresponding Weyl group and $w_0 \in W$ be the longest element with the length l_0 .

In [11, Proposition 4.2] we have shown by using the braid-type isomorphisms that $B(\lambda)$ can be embedded in the finite rank $\mathbf{Z}$ -lattice $\mathbf{Z}^{l_0}$. Here we obtain its simpler proof as an application of Proposition 3.1. Indeed, in this case, since $V(\lambda) = V_{w_0}(\lambda)$, we have $B(\lambda) = B_{w_0}(\lambda)$. This implies:

Proposition 3.4 *There exists the following embedding,*

$$\Psi_\iota^{(\lambda)} : B(\lambda) (= B_{w_0}(\lambda)) \xrightarrow{\sim} \Sigma_{w_0}[\lambda] \hookrightarrow \mathbf{Z}^{l_0}, \quad (3.19)$$

where ι is an infinite sequence of indices such that its first l_0 subsequence $i_{l_0}, i_{l_0-1}, \cdots, i_1$ is a reduced longest word associated with the longest element w_0 (see [11, 4.2]).

4 Extremal vectors

We still keep the notations of 3.2 and we do not necessarily assume that (λ, ι) is ample.

4.1 Explicit description of extremal vectors

For $w \in W$, we call $w\lambda$ the extremal weight of $B(\lambda)$ and call the unique element $u_{w\lambda} \in B(\lambda)_{w\lambda}$ *extremal vector* with the extremal weight $w\lambda$.

The image of $u_{w\lambda}$ by $\Psi_t^{(\lambda)}$ is included in $\Sigma_w[\lambda]$. We are going to determine it by the following way.

Proposition 4.1 *For $w \in W$ ($\text{length}(w) = L$), set $x_w = (\cdots, x_k, \cdots, x_2, x_1) := \Psi_t^{(\lambda)}(u_{w\lambda})$. Then the element x_w is given as the unique solution of the following system of linear equations:*

$$\begin{cases} x_k = 0 & \text{for } k > L, \\ \beta_k^{(-)}(x) = 0 & \text{for } k \leq L, \end{cases} \quad (4.1)$$

where the linear function $\beta_k^{(-)}$ is as in (2.11).

Proof. The equations $\text{eq}(L)$

$$\beta_1^{(-)}(x) = \beta_2^{(-)}(x) = \cdots = \beta_L^{(-)}(x) = 0.$$

is the system of the linear equations in indeterminates $x_1, x_2, \cdots, x_L$. If we write $\text{eq}(L)$ in a matrix form $A\vec{x} = \vec{\xi}$ where $\vec{x} = {}^t(x_1, \cdots, x_L)$, due to the explicit form of $\beta_k^{(-)}$ in (2.11), the matrix A is a triangular integer matrix whose diagonal entries are all 1 and the vector $\vec{\xi} = {}^t(\xi_1, \cdots, \xi_L)$ is given by $\xi_k = \langle h_{i_k}, \lambda \rangle$ if $k^{(-)} = 0$ and otherwise $\xi_k = 0$. Thus, the equation $\text{eq}(L)$ can be solved uniquely and all the entries are integers. We set the solution $(y_1, \cdots, y_L)$. Therefore, it suffices to show $x_w := \Psi_t^{(\lambda)}(u_{w\lambda}) = (\cdots, 0, 0, y_L, \cdots, y_1)$. Let us show this by the induction on the length of w . Set $w := s_{i_L} s_{i_{L-1}} \cdots s_{i_2} s_{i_1}$, $w' := s_{i_{L-1}} \cdots s_{i_2} s_{i_1}$, $y_w := (\cdots, 0, 0, y_L, y_{L-1}, \cdots, y_2, y_1)$ and $y_{w'} := (\cdots, 0, 0, y_{L-1}, \cdots, y_2, y_1)$. Note that $y_{w'}$ is the unique solution of $\text{eq}(L-1)$ and also the image of $x_{w'}$ by $\Psi_t^{(\lambda)}$ from the hypothesis of the induction. Here we show the following lemma:

Lemma 4.2 *For w and w' as above, let $u_{w\lambda}$ and $u_{w'\lambda}$ be the corresponding extremal vectors. Then we have*

$$u_{w\lambda} = \tilde{f}_{i_L}^{\max} u_{w'\lambda}, \quad (4.2)$$

where $\tilde{f}_i^{\max} u := \tilde{f}_i^{\varphi_i(u)} u$.

Proof of Lemma 4.2. By the definition of $\tilde{f}_i^{\max}$, $\tilde{f}_{i_L}^{\max} u_{w'\lambda} \neq 0$. Owing to the uniqueness of the extremal vector, it suffices to show

$$wt(\tilde{f}_{i_L}^{\max} u_{w'\lambda}) = w\lambda. \quad (4.3)$$

Let S_L be the i_L -string in $B(\lambda)$ including $u_{w'\lambda}$. By Theorem 1.1 (iii), we know that $S_L \cap B_{w'}(\lambda)$ is equal to (1) S_L or (2) {highest weight vector in S_L }. In the case (1), $u_{w'\lambda}$ is the lowest weight vector in S_L since $\mu \succ w'\lambda$ when $B_{w'}(\lambda)_\mu \neq \emptyset$. This implies $\langle h_{i_L}, w'\lambda \rangle \leq 0$. Suppose that $\langle h_{i_L}, w'\lambda \rangle < 0$. Then we have

$$w\lambda = s_{i_L}(w'\lambda) = w'\lambda - \langle h_{i_L}, w'\lambda \rangle \alpha_{i_L} \succ w'\lambda, \quad (4.4)$$

which contradicts Theorem 1.1 (ii). Thus, in this case we have $\langle h_{i_L}, w'\lambda \rangle = 0$ and then the length of $S_L = 0$. This means $\varphi_{i_L}(u_{w'\lambda}) = 0$ and then $u_{w\lambda} = u_{w'\lambda}$. In the case (2), since $\varepsilon_{i_L}(u_{w'\lambda}) = 0$, we have $\varphi_{i_L}(u_{w'\lambda}) = \langle h_{i_L}, wt(u_{w'\lambda}) \rangle$ and then

$$\begin{aligned} wt(\tilde{f}_{i_L}^{\max} u_{w'\lambda}) &= w'\lambda - \varphi_{i_L}(u_{w'\lambda}) \alpha_{i_L} \\ &= w'\lambda - \langle h_{i_L}, wt(u_{w'\lambda}) \rangle \alpha_{i_L} \\ &= s_{i_L}(w'\lambda) = w\lambda. \end{aligned}$$

Now, we obtain (4.3) and then completed the proof of Lemma 4.2 $\square$

By this lemma, it suffices to show

$$\tilde{f}_{i_L}^{\max} y_{w'} = y_w, \quad (4.5)$$

Let us see how $\tilde{f}_{i_L}$ acts on $y_{w'}$. For $k \in \mathbf{Z}_{\geq 1}$ and $m \in \mathbf{Z}_{>0}$ we set $k^{(\pm 1)} := k^{(\pm)}$, $k^{(+m)} := (k^{(+m-1)})^{(+)}$ and $k^{(-m)} := (k^{(-m-1)})^{(-)}$ (as for $k^{(\pm)}$, see 2.3). For $m \geq 1$, we have

$$\sigma_{L(-m)} + \beta_{L(-m)}^{(-)} = \sigma_{L(-m-1)}. \quad (4.6)$$

Since $\beta_k^{(-)}(y_{w'}) = 0$ for $k \leq L-1$, we have by (4.6)

$$\sigma_{L(-)}(y_{w'}) = \sigma_{L(-2)}(y_{w'}) = \cdots = \sigma_{L(-m)}(y_{w'}) = \cdots = \sigma_0^{(i_L)}(y_{w'}). \quad (4.7)$$

Now we consider the following cases:

$$(a) \beta_L^{(-)}(y_{w'}) < 0. \quad (b) \beta_L^{(-)}(y_{w'}) \geq 0.$$

In the case (a), we have

$$\begin{aligned} 0 &= \cdots = \sigma_{L(+m)}(y_{w'}) = \cdots = \sigma_L(y_{w'}) \\ &> \sigma_{L(-)}(y_{w'}) = \cdots = \sigma_{L(-m)}(y_{w'}) = \cdots = \sigma_0^{(i_L)}(y_{w'}). \end{aligned} \quad (4.8)$$

It follows from (3.3) that

$$\tilde{f}_{i_L}(y_{w'}) = (\cdots, 1, y_{L-1}, \cdots, y_1). \quad (4.9)$$

In the case (b), we have

$$\begin{aligned} 0 &= \cdots = \sigma_{L(+m)}(y_{w'}) = \cdots = \sigma_L(y_{w'}) \\ &\leq \sigma_{L(-)}(y_{w'}) = \cdots = \sigma_{L(-m)}(y_{w'}) = \cdots = \sigma_0^{(i_L)}(y_{w'}), \end{aligned} \quad (4.10)$$

which implies $\tilde{f}_{i_L}(y_{w'}) = 0$ by (3.3). In this case, by (3.5) and (3.6) we have

$$\varepsilon_{i_L}(y_{w'}) = \sigma_0^{(i_L)}(y_{w'}) = -\langle h_{i_L}, wt(y_{w'}) \rangle, \quad (4.11)$$

and then also by (3.7) we have

$$\varphi_{i_L}(y_{w'}) = \varepsilon_{i_L}(y_{w'}) + \langle h_{i_L}, wt(y_{w'}) \rangle = 0, \quad (4.12)$$

which implies that $y_{w'}$ is the lowest weight vector in the i_L -string including $y_{w'}$. Thus, the case (b) corresponds to the case (1) in the proof of Lemma 4.2. So the length of i_L -string is 0 and then we have $\varepsilon_{i_L}(y_{w'}) = 0$. By (4.10) and (4.11) we have $0 = \sigma_0^{(i_L)}(y_{w'}) = \cdots = \sigma_{L(-)}(y_{w'}) = \sigma_L(y_{w'}) = 0$, and then $\beta_L^{(-)}(y_{w'}) = 0$ by (4.6), which means $y_w = y_{w'} (= \tilde{f}_{i_L}^{\max} y_{w'})$, that is, (4.5) with $y_L = 0$.

In the case (a), we can suppose that $\varphi_{i_L}(y_{w'}) > 0$. Let us show

$$\tilde{f}_{i_L}^k(y_{w'}) = (\cdots, 0, 0, k, y_{L-1}, \cdots, y_2, y_1), \quad (4.13)$$

for $1 \leq k \leq \varphi_{i_L}(y_{w'})$ by the induction on k . Assuming (4.13) ($1 \leq k < \varphi_{i_L}(y_{w'})$), let us see $\tilde{f}_{i_L}^{k+1}(y_{w'})$. Set $\bar{y} := \tilde{f}_{i_L}^k(y_{w'}) = (\cdots, 0, 0, k, y_{L-1}, \cdots, y_2, y_1)$. In the case (a), by the argument for the case (1) in the proof of Lemma 4.2, we have $\varepsilon_{i_L}(y_{w'}) = 0$ and then by $\sigma_L(y_{w'}) = 0$ and (3.8),

$$\varphi_{i_L}(y_{w'}) = \langle h_{i_L}, wt(y_{w'}) \rangle = -\sigma_0^{(i_L)}(y_{w'}) = \cdots = -\sigma_{L(-)}(y_{w'}) = -\beta_L^{(-)}(y_{w'}) \quad (4.14)$$

Set $F := \varphi_{i_L}(y_{w'}) = -\beta_L^{(-)}(y_{w'})$. On the other hand, if $i_l = i_L$, we have

$$\sigma_l(\bar{y}) = \sigma_l(y_{w'}) + 2k = -F + 2k \quad (l < L), \quad \sigma_L(\bar{y}) = k. \quad (4.15)$$

It follows from (4.14) and (4.15) that if $k < F$, we have $\sigma_{L(-n)}(\bar{y}) = 2k - F < k = \sigma_L(\bar{y}) \geq 0 = \sigma_{L(+m)}(\bar{y})$ ($m, n > 0$) and then by (3.3)

$$\tilde{f}_{i_L}(\bar{y}) = (\cdots, 0, 0, k+1, y_{L-1}, \cdots, y_1). \quad (4.16)$$

Hence, we obtain

$$\tilde{f}_{i_L}^{\max}(y_{w'}) = \tilde{f}_{i_L}^F(y_{w'}) = (\cdots, 0, 0, -\beta_L^{(-)}(y_{w'}), y_{L-1}, \cdots, y_1). \quad (4.17)$$

Here $(y_1, \cdots, y_{L-1}, -\beta_L^{(-)}(y_{w'}))$ satisfies the equations eq(L) and then it follows that $y_w = \tilde{f}_{i_L}^{\max}(y_{w'})$. $\square$

Remark. Note that we obtain Proposition 4.1 without the assumption “ample”. In [11, Example 3.9], we introduced the “non-ample” example: $\mathfrak{g} = A_3$ and $\iota = 212321$. But, even in this case, applying Proposition 4.1 we have

$$\begin{aligned} x_{s_3 s_2 s_1} &= (0, 0, 0, \lambda_1 + \lambda_2 + \lambda_3, \lambda_1 + \lambda_2, \lambda_1), \\ x_{s_2 s_3 s_2 s_1} &= (0, 0, \lambda_3, \lambda_1 + \lambda_2 + \lambda_3, \lambda_1 + \lambda_2, \lambda_1), \\ x_{s_1 s_2 s_3 s_2 s_1} &= (0, \lambda_2 + \lambda_3, \lambda_3, \lambda_1 + \lambda_2 + \lambda_3, \lambda_1 + \lambda_2, \lambda_1), \\ x_{s_2 s_1 s_2 s_3 s_2 s_1} &= (\lambda_2, \lambda_2 + \lambda_3, \lambda_3, \lambda_1 + \lambda_2 + \lambda_3, \lambda_1 + \lambda_2, \lambda_1), \end{aligned}$$

where $\lambda_i = \langle h_i, \lambda \rangle$.

4.2 Rank 2 cases

We apply Proposition 4.1 to arbitrary rank 2 cases.

First we review the result in [10]. The setting here is same as those in [10]. We set $I = \{1, 2\}$, and $\iota = (\cdots, 2, 1, 2, 1)$. The Cartan matrix is given by:

$$\langle h_1, \alpha_1 \rangle = \langle h_2, \alpha_2 \rangle = 2, \quad \langle h_1, \alpha_2 \rangle = -c_1, \quad \langle h_2, \alpha_1 \rangle = -c_2.$$

Here we either have $c_1 = c_2 = 0$, or both c_1 and c_2 are positive integers. We set $X = c_1 c_2 - 2$, and define the integer sequence $a_l = a_l(c_1, c_2)$ for $l \geq 0$ by setting $a_0 = 0$, $a_1 = 1$ and, for $k \geq 1$,

$$a_{2k} = c_1 P_{k-1}(X), \quad a_{2k+1} = P_k(X) + P_{k-1}(X), \quad (4.18)$$

where the $P_k(X)$ are *Chebyshev polynomials* given by the following generating function:

$$\sum_{k \geq 0} P_k(X) z^k = (1 - Xz + z^2)^{-1}. \quad (4.19)$$

Here define $a'_l(c_1, c_2) := a_l(c_2, c_1)$. The several first Chebyshev polynomials and terms a_l are given by

$$P_0(X) = 1, \quad P_1(X) = X, \quad P_2(X) = X^2 - 1, \quad P_3(X) = X^3 - 2X,$$

$$a_2 = c_1, \quad a_3 = c_1 c_2 - 1, \quad a_4 = c_1(c_1 c_2 - 2),$$

$$a_5 = (c_1 c_2 - 1)(c_1 c_2 - 2) - 1, \quad a_6 = c_1(c_1 c_2 - 1)(c_1 c_2 - 3).$$

Let $l_{\max} = l_{\max}(c_1, c_2)$ be the minimal index l such that $a_{l+1} < 0$ (if $a_l \geq 0$ for all $l \geq 0$, then we set $l_{\max} = +\infty$). By inspection, if $c_1 c_2 = 0$ (resp. $1, 2, 3$) then $l_{\max} = 2$ (resp. $3, 4, 6$). Furthermore, if $c_1 c_2 \leq 3$ then $a_{l_{\max}} = 0$ and $a_l > 0$ for $1 \leq l < l_{\max}$. On the other hand, if $c_1 c_2 \geq 4$, i.e., $X \geq 2$, it is easy to see from (4.19) that $P_k(X) > 0$ for $k \geq 0$, hence $a_l > 0$ for $l \geq 1$; in particular, in this case $l_{\max} = +\infty$.

Proposition 4.3 ([10]) *In the rank 2 case, for a dominant integral weight $\lambda = m_1 \Lambda_1 + m_2 \Lambda_2$ ($m_1, m_2 \in \mathbf{Z}_{\geq 0}$) the image of the embedding $\Psi_t^{(\lambda)}$ is given by*

$$\text{Im}(\Psi_t^{(\lambda)}) = \left\{ (\cdots, x_2, x_1) \in \mathbf{Z}_{\geq 0}^\infty : \begin{array}{l} x_k = 0 \text{ for } k > l_{\max}, \quad m_1 \geq x_1, \\ a_l x_l - a_{l-1} x_{l+1} \geq 0, \\ m_2 + a'_{l+1} x_l - a'_l x_{l+1} \geq 0, \\ \text{for } 1 \leq l < l_{\max} \end{array} \right\}. \quad (4.20)$$

Note that the cases when $l_{\max} < +\infty$, or equivalently, the image $\text{Im}(\Psi_t^{(\lambda)})$ is contained in a lattice of finite rank, just correspond to the Lie algebras $\mathfrak{g} = A_1 \times A_1, A_2, B_2$ or C_2, G_2 .

For $L \in \mathbf{Z}_{\geq 0}$ ($L \leq l_{\max}$), we set

$$w_L := \begin{cases} s_1(s_2 s_1)^l & \text{if } L = 2l + 1, \\ (s_2 s_1)^l & \text{if } L = 2l. \end{cases} \quad (4.21)$$

For a dominant integrable weight $\lambda = m_1 \Lambda_1 + m_2 \Lambda_2 \in P_+$, we define the integer sequence $d_k = d_k(c_1, c_2)$ ($k \geq 1$) as follows:

$$d_k(c_1, c_2) := m_1 a_k(c_2, c_1) + m_2 a_{k-1}(c_1, c_2), \quad (4.22)$$

where $\{a_k\}_{k \geq 0}$ is the integer sequence given in (4.18).

Proposition 4.4 *The image x_{w_L} of the extremal vector $u_{w_L\lambda} \in B_{w_L}(\lambda)$ ($\lambda = m_1\Lambda_1 + m_2\Lambda_2 \in P_+$) associated with $w_L \in W$ can be described as follows:*

$$x_{w_L} \left(:= \Psi_l^{(\lambda)}(u_{w_L\lambda}) \right) = (\cdots, 0, 0, d_L, d_{L-1}, \cdots, d_2, d_1). \quad (4.23)$$

Proof. In this setting, we have

$$\begin{aligned} \beta_1^{(-)}(x) &= x_1 - m_1, \quad \beta_2^{(-)}(x) = x_2 - c_2x_1 - m_2, \\ \beta_{2k+1}^{(-)}(x) &= x_{2k+1} - c_1x_{2k} + x_{2k-1}, \quad \beta_{2k+2}^{(-)}(x) = x_{2k+2} - c_2x_{2k+1} + x_{2k}, \quad (k \geq 1). \end{aligned}$$

We shall show $d_1, d_2, \cdots, d_L$ are the solutions of the equations

$$\beta_1^{(-)}(x) = \beta_2^{(-)}(x) = \cdots = \beta_L^{(-)}(x) = 0. \quad (4.24)$$

Solving $\beta_1^{(-)}(x) = \beta_2^{(-)}(x) = 0$, we have

$$x_1 = m_1 = m_1a'_1 + m_2a_0 = d_1, \quad x_2 = m_1c_2 + m_2 = m_1a'_2 + m_2a_1 = d_2. \quad (4.25)$$

Here note that we can write $d_k = m_1a'_k + m_2a_{k-1}$. By the definition of a_k , we can easily see that $\{a_k\}$ (resp. $\{a'_k\}$) is uniquely determined by

$$\begin{aligned} a_0 &= 0, \quad a_1 = 1, \quad a_{2k+1} = c_2a_{2k} - a_{2k-1}, \quad a_{2k+2} = c_1a_{2k+1} - a_{2k}, \quad (k \geq 0). \\ (\text{resp. } a'_0 &= 0, \quad a'_1 = 1, \quad a'_{2k+1} = c_1a'_{2k} - a'_{2k-1}, \quad a'_{2k+2} = c_2a'_{2k+1} - a'_{2k}). \end{aligned}$$

Here for $k \geq 1$ we have $d_{2k+1} - c_1d_{2k} + d_{2k-1} = m_1(a'_{2k+1} - c_1a'_{2k} + a'_{2k-1}) + m_2(a_{2k} - c_1a_{2k-1} + a_{2k-2}) = 0$ and $d_{2k+2} - c_2d_{2k+1} + d_{2k} = m_1(a'_{2k+2} - c_2a'_{2k+1} + a'_{2k}) + m_2(a_{2k+1} - c_2a_{2k} + a_{2k-1}) = 0$, which implies that $(d_1, d_2, \cdots, d_L)$ is the unique solution of (4.24). Now, we obtain the desired result. $\square$

In conclusion of this section, we illustate the case of $A_1^{(1)}$, that is, $c_1 = c_2 = 2$. In this case, $X = c_1c_2 - 2 = 2$. It follows at once from (4.19) that $P_k(2) = k + 1$; hence, (4.18) gives $a_l = l$ for $l \geq 0$. We see that for type $A_1^{(1)}$,

$$x_{w_L} = (\cdots, 0, 0, Lm_1 + (L-1)m_2, \cdots, km_1 + (k-1)m_2, \cdots, 2m_1 + m_2, m_1).$$

Acknowledgements The author would like to acknowledge M.Okado for his interest to my work. Indeed, this work was motivated by his question about the relation of polyhedral realizations and the crystals of Demazure modules.

References

- [1] Kashiwara M, Crystallizing the q -analogue of universal enveloping algebras, *Commun.Math.Phys.*, **133** (1990), 249–260.
- [2] Kashiwara M, On crystal bases of the q -analogue of universal enveloping algebras, *Duke Math. J.*, **63** (1991) 465–516.
- [3] Kashiwara M, Crystal base and Littelmann’s refined Demazure character formula, *Duke Math. J.*, **71**(1993), 839–858.

- [4] Kashiwara M, Crystallization of quantized enveloping algebras, Sugaku Expositions, **7**, No.1, June, 1994.
- [5] Kashiwara M and Nakashima T, Crystal graph for representations of the q -analogue of classical Lie algebras, *J. Algebra*, **165**, (1994), 295–345.
- [6] S.Kumar, Demazure character formula in arbitrary Kac-Moody setting, *Invent. Math.*, **89** (1987), 395–423.
- [7] Littelmann P, Crystal Graphs and Young Tableaux, *J.Algebra*, **175**, 65–87 (1995).
- [8] Mathieu O., Formules de caractères pour les algèbres de Kac-Moody générales, *Astérisue* **159–160**, (1988).
- [9] Nakashima T and Zelevinsky A, Polyhedral Realization of Crystal Bases for Quantized kac-Moody Algebras, *Advances in Mathematics*, **131**, No.1, (1997), 253–278.
- [10] Nakashima T., Polyhedral Realizations of Crystal Bases for Integrable Highest Weight Modules, *J.Algebra* **219**, 571–597 (1999).
- [11] Nakashima T., Polyhedral Realizations of Crystal Bases and Braid-type Isomorphisms, preprint, mathQA/9906100, (to appear in Contemporary Mathematics).